

Certifying the Intersection of Reach Sets of Integrator Agents with Set-valued Input Uncertainties

Shadi Haddad and Abhishek Halder

Department of Applied Mathematics
University of California, Santa Cruz


Conference on Decision and Control, Dec 8, 2022

Reach Set



$$\mathcal{X}_t := \{ \mathbf{x}(t) \in \mathbb{R}^n \mid \dot{\mathbf{x}} = f(\mathbf{x}) + g(\mathbf{x})\mathbf{u},$$

$$\mathbf{x}(t=0) = \mathbf{x}_0, \mathbf{u}(s) \in \mathcal{U}(s)$$



for all $0 \leq s \leq t\}$




Safety critical applications such as motion planning & collision warning systems



Existing Algorithms



[Cohen et. al, 1995]



[Elmer et. al, 1988]

No specific algebraic results about the ground truth

Difficult to quantitatively compare performance

Our approach

Generic → specific algorithm exploiting geometry of the true set

Contribution

Integrator reach set

Brunovsky normal form

$$\mathcal{X}_t := \{\mathbf{x}(t) \in \mathbb{R}^n \mid \mathbf{Ax} + \mathbf{Bu}, \mathbf{x}(t=0) = x_0, \mathbf{u}(s) \in \mathcal{U}(s), \forall 0 \leq s \leq t\}$$

$$\mathcal{U}(s) := \{\mathbf{u}(s) \in \mathbb{R}^m \mid \|\mathbf{u}(s)\|_p \leq \ell(s)\}, \forall s \in [0, t]$$

$$\mathbf{A} := \text{blkdiag}(\mathbf{A}_1, \dots, \mathbf{A}_m), \quad \mathbf{B} := \text{blkdiag}(\mathbf{b}_1, \dots, \mathbf{b}_m),$$

$$\mathbf{A}_j = \begin{bmatrix} \mathbf{0} & \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 & \dots & \mathbf{e}_{r_j-1} \end{bmatrix}, \quad \mathbf{b}_j = \mathbf{e}_{r_j}$$

$$\mathbf{r} = (r_1, r_2, \dots, r_m)^\top \in \mathbb{Z}_+^m \quad \text{Relative degree vector}$$

Benchmarking the performance of over-approximation algorithms

Predicting the intersection between differentially flat nonlinear systems

Motivation

Dynamics of VTOL aircraft

$$\dot{z}_1 = z_2$$

$$\dot{z}_2 = -v_1 \sin(z_5) + \epsilon v_2 \cos(z_5)$$

$$\dot{z}_3 = z_4$$


$$\dot{z}_4 = v_1 \cos(z_5) + \epsilon v_2 \sin(z_5) - g$$

$$\dot{z}_5 = z_6$$

$$\dot{z}_6 = v_2$$

Normal form

$$\dot{x} = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} x + \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ v_2 \end{pmatrix}$$



The VTOL aircraft

Intersection in normal coordinate x




Intersection in original coordinates z

Integrator reach sets are in general, compact and convex

Intersection Detection

General formula for the collision detection

$$\text{dist}(A, B) := \min_{\mathbf{x}^A \in \mathcal{X}_t^A, \mathbf{x}^B \in \mathcal{X}_t^B} \|\mathbf{x}^A - \mathbf{x}^B\|_2^2$$




Lack of analytical handle on the boundary!!!

Alternative approach, intersection occurs if

$$\mathbf{0} \in \{\mathcal{X}_t^A - \mathcal{X}_t^B\}$$

Minkowski Difference



Support function $h_{\mathcal{X}_t}(\mathbf{y}) = \max_{\mathbf{x} \in \mathcal{X}} \langle \mathbf{x}^\top, \mathbf{y} \rangle$

$$\mathcal{X}_t^A \cap \mathcal{X}_t^B \neq \emptyset \Leftrightarrow \mathbf{0} \in \mathcal{X}_t^A - \mathcal{X}_t^B \Leftrightarrow$$

$$\forall \mathbf{y} \in \mathbb{S}^{n-1}, h_{\mathcal{X}_t^A - \mathcal{X}_t^B}(\mathbf{y}) \geq 0$$

Intersection Detection

Intersection Detection Oracle

$$\min_{\mathbf{y} \in \mathbb{S}^{n-1}} \{h_{\mathcal{X}_t^A}(\mathbf{y}) + h_{\mathcal{X}_t^B}(-\mathbf{y})\} \geq (<) 0 \iff \mathcal{X}_t^A \cap \mathcal{X}_t^B \neq (=) \emptyset$$

Theorem: The support function of \mathcal{X}_t

$$h_{\mathcal{X}_t}(\mathbf{y}) = \sum_{j=1}^m \langle \mathbf{y}_j, \exp(t\mathbf{A})\mathbf{x}_{j0} \rangle + \int_0^t \ell(s) \|(\exp(s\mathbf{A})\mathbf{B})^\top \mathbf{y}\|_q ds, \quad 1/p + 1/q = 1$$

Discretize $[0, t]$ into K intervals:

$$\int_0^t \|(\exp(s\mathbf{A})\mathbf{B})^\top \mathbf{y}\|_q ds \approx \cdot$$

$$\frac{\Delta s}{2} \sum_{k=1}^K \left(\|(\exp(s_{k-1}\mathbf{A})\mathbf{B})^\top \mathbf{y}\|_q + \|(\exp(s_k\mathbf{A})\mathbf{B}(s_k))^\top \mathbf{y}\|_q \right)$$


Lossless Convexification

Nonconvex Optimization Problem

$$\overbrace{p^* = \min_{\boldsymbol{\eta} \in \mathbb{R}^{n+K+1}} \langle \boldsymbol{\kappa}(t), \boldsymbol{\eta} \rangle}^{\text{Nonconvex Objective}}$$

s. $\|M_k \boldsymbol{\eta}\|_q - \langle \mathbf{e}_{n+k}^{n+K+1}, \boldsymbol{\eta} \rangle \leq 0,$

$-\tilde{\mathbf{N}}\boldsymbol{\eta} \leq \mathbf{0}, \quad \|\mathbf{N}\boldsymbol{\eta}\|_2 \leq 1$



Nonconvex Constraint

→
Lossless
Convexification

Second-Order Cone Program

$$\overbrace{\tilde{p}^* = \min_{\boldsymbol{\eta} \in \mathbb{R}^{n+K+1}} \langle \boldsymbol{\kappa}(t), \boldsymbol{\eta} \rangle}^{\text{Convex Objective}}$$

s. $\|M_k \boldsymbol{\eta}\|_q - \langle \mathbf{e}_{n+k}^{n+K+1}, \boldsymbol{\eta} \rangle \leq 0,$

$-\tilde{\mathbf{N}}\boldsymbol{\eta} \leq \mathbf{0}, \quad \|\mathbf{N}\boldsymbol{\eta}\|_2 \leq 1$

(i) $\tilde{p}^* \leq 0$

(ii) $\tilde{p}^* = 0 \Rightarrow 0 \leq p^* \Leftrightarrow \mathcal{X}_t^A \cap \mathcal{X}_t^B \neq \emptyset$

(iii) $\tilde{p}^* < 0 \Rightarrow \tilde{p}^* = p^* < 0 \Leftrightarrow \mathcal{X}_t^A \cap \mathcal{X}_t^B = \emptyset$


Example

$$\dot{\mathbf{x}}^i = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \mathbf{x}^i + \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \\ \hline 0 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{u}^i, \quad i \in \{\text{A, B}\}, \quad r = (3, 2)^\top$$

∞ -norm ball


$$\mathcal{U}(s) := [\alpha_1(s), \beta_1(s)] \times [\alpha_2(s), \beta_2(s)] \quad \text{for all } 0 \leq s \leq t$$

Decoupled dynamics for each block



$$\begin{aligned} \mathcal{X}_t^i &= \mathcal{X}_{1t}^i \times \mathcal{X}_{2t}^i = \mathcal{X}_{1t}^i + \mathcal{X}_{2t}^i \\ h_{\mathcal{X}}^i &= h_{\mathcal{X}_{1t}}^i + h_{\mathcal{X}_{2t}}^i \quad i \in \{\text{A, B}\} \end{aligned}$$

Example



$$(\tilde{p}_1^*, \tilde{p}_2^*) = (0, -0.54)$$


Computation time ≈ 1.24 s

$$\mathcal{X}_{1t}^A \cap \mathcal{X}_{1t}^B \neq \emptyset \quad \text{and} \quad \mathcal{X}_{2t}^A \cap \mathcal{X}_{2t}^B = \emptyset \iff \boxed{\mathcal{X}_t^A \cap \mathcal{X}_t^B = \emptyset}$$


Ongoing Work

Map them back to original coordinate
 z via known diffeomorphism

Compute the reach set and its
functionals in normal coordinate x



$$\text{vol}(\mathcal{Z}_t) = 206.7362$$



$$\text{vol}(\mathcal{X}_t) = 15.4292$$

Thank You