



National
Science
Foundation

Schrödinger Bridges over Kinetic Swarming Models

Asmaa Eldesoukey*, Md Zulfikur Haider**, Italo Napolitano***, Yongxin Chen****, Abhishek Halder*

*Department of Aerospace Engineering, Iowa State University, Ames, Iowa 50010, USA.


**Department of Mathematics, Iowa State University, Ames, Iowa 50010, USA.

***Modeling and Engineering Risk and Complexity Department, Scuola Superiore Meridionale, via Mezzocannone 4, 80134, Naples, Italy.

****School of Aerospace Engineering, Georgia Institute of Technology, Atlanta, GA 30332, USA.



SCHRÖDINGER BRIDGES



KINETIC SWARM MODELS

CUCKER-SMALE KINETICS

$$\begin{cases} dX_t^i = V_t^i dt, \\ dV_t^i = \frac{1}{N} \sum_{j \neq i, j=1}^N a(\|X_t^i - X_t^j\|) (V_t^j - V_t^i) dt + \sigma dB_t^i, \end{cases} \quad \begin{cases} dX_t^i = V_t^i dt, \\ dV_t^i = -\frac{1}{N} \sum_{j \neq i} \nabla_{X_t^i} W(\|X_t^i - X_t^j\|) dt + \sigma dB_t^i, \end{cases}$$

$$a(z) := \frac{K}{(1+z^2)^\gamma}, \quad K > 0, \gamma \geq 0. \quad W(z) := C_R e^{-z/\ell_R} - C_A e^{-z/\ell_A},$$

MORSE KINETICS

STOCHASTIC OPTIMAL CONTROL PROBLEM

$$\text{Minimize}_{u, \mu} \quad \frac{1}{2\sigma^2} \int_0^T \int_{\mathbb{R}^{2d}} \|u_t(x, v)\|^2 \mu_t dx dv dt,$$

$$\text{subject to} \quad \partial_t \mu_t + \langle v, \nabla_x \mu_t \rangle + \nabla_v \cdot (\mu_t (\mathcal{I}_t[\rho_t] + u_t)) = \frac{\sigma^2}{2} \Delta_v \mu_t,$$

$$\text{with} \quad \mu_0(x, v) = \mu_{\text{initial}}(x, v), \quad \mu_T(x, v) = \mu_{\text{final}}(x, v).$$

SCHRÖDINGER SYSTEM: CUCKER-SMALE KINETICS

$$\begin{cases} \mu_t^*(x, v) = \varphi_t(x, v) \hat{\varphi}_t(x, v), \\ \partial_t \varphi_t + \langle v, \nabla_x \varphi_t \rangle + \langle f_t[\mu_t^*], \nabla_v \varphi_t \rangle + \frac{\sigma^2}{2} \Delta_v \varphi_t = r_t(x, v) \varphi_t, \\ \partial_t \hat{\varphi}_t + \langle v, \nabla_x \hat{\varphi}_t \rangle + \nabla_v \cdot (\hat{\varphi}_t f_t[\mu_t^*]) - \frac{\sigma^2}{2} \Delta_v \hat{\varphi}_t = -r_t(x, v) \hat{\varphi}_t, \end{cases}$$

$$\text{where} \quad f_t[\mu_t^*](x, v) = \int_{\mathbb{R}^{2d}} a(\|x - \tilde{x}\|) \mu_t^*(\tilde{x}, \tilde{v}) (\tilde{v} - v) d\tilde{x} d\tilde{v},$$

$$r_t(x, v) = \int_{\mathbb{R}^{2d}} a(\|x - \tilde{x}\|) \hat{\varphi}_t(\tilde{x}, \tilde{v}) \langle \tilde{v} - v, \nabla_{\tilde{v}} \varphi_t(\tilde{x}, \tilde{v}) \rangle d\tilde{x} d\tilde{v},$$

subject to

$$\varphi_0(x, v) \hat{\varphi}_0(x, v) = \mu_{\text{initial}}(x, v),$$

$$\varphi_T(x, v) \hat{\varphi}_T(x, v) = \mu_{\text{final}}(x, v).$$

SCHRÖDINGER SYSTEM: MORSE KINETICS

$$\begin{cases} \mu_t^*(x, v) = \varphi_t(x, v) \hat{\varphi}_t(x, v), \\ \partial_t \varphi_t + \langle v, \nabla_x \varphi_t \rangle + \langle g_t[\rho_t^*], \nabla_v \varphi_t \rangle + \frac{\sigma^2}{2} \Delta_v \varphi_t = s_t(x) \varphi_t, \\ \partial_t \hat{\varphi}_t + \langle v, \nabla_x \hat{\varphi}_t \rangle + \langle g_t[\rho_t^*], \nabla_v \hat{\varphi}_t \rangle - \frac{\sigma^2}{2} \Delta_v \hat{\varphi}_t = -s_t(x) \hat{\varphi}_t, \end{cases}$$

$$\text{where} \quad g_t[\rho_t^*](x) = - \int_{\mathbb{R}^d} \nabla_x W(\|x - \tilde{x}\|) \rho_t^*(\tilde{x}) d\tilde{x},$$




$$s_t(x) := - \int_{\mathbb{R}^{2d}} \langle \nabla_x W(\|x - \tilde{x}\|), \nabla_{\tilde{v}} \varphi_t(\tilde{x}, \tilde{v}) \rangle \hat{\varphi}_t(\tilde{x}, \tilde{v}) d\tilde{x} d\tilde{v},$$

subject to



$$\varphi_0(x, v) \hat{\varphi}_0(x, v) = \mu_{\text{initial}}(x, v),$$

$$\varphi_T(x, v) \hat{\varphi}_T(x, v) = \mu_{\text{final}}(x, v).$$


CUCKER-SMALE MFSB



MORSE MFSB



MFSB WITH PARTIAL STATE INFORMATION



REFERENCES

- [1] F. Cucker and E. Mordecki, "Flocking in noisy environments", *Journal de mathématiques pures et appliquées* 89, 278, 2008.
- [2] F. Cucker and S. Smale, "Emergent behavior in flocks", *IEEE Transactions on Automatic Control* 52, 852, 2007.
- [3] F. Cucker and S. Smale, "On the mathematics of emergence", *Japanese Journal of Mathematics* 2, 197, 2007.
- [4] A. Mogilner, L. Edelstein-Keshet, L. Gent, and A. Spiros, "Mutual interactions, potentials, and individual distance in a social aggregation", *Journal of mathematical biology* 47, 353, 2003.
- [5] A. Eldesoukey, M. Z. Haider, I. Napolitano, Y. Chen, and A. Halder, "Schrödinger bridges over kinetic swarming models," *arXiv preprint, arXiv:2608.25281*, 2026.

The authors gratefully acknowledge the supports of the NSF grants: 2111688, 2450377, 2450378.